



The Projective Rigidity of Projective Space


Michael Eastwood

[joint work with Laurent Stolovitch]

Australian National University

Blaschke conjecture/theorem

On a sphere



- all geodesics are closed
- all geodesics have the same length

The same features are present (in any dimension) on

- real projective space
- complex projective space

CROSSes

Blaschke rigidity

Deformations

- Riemannian $g_{ab} \mapsto \tilde{g}_{ab} = g_{ab} + \epsilon h_{ab}$
- Projective $\nabla_a \mapsto \tilde{\nabla}_a = \nabla_a + \epsilon \Gamma_a$

Two-sphere with round metric g_{ab}

WLG $\tilde{g}_{ab} = (1 + \epsilon f)^2 g_{ab}$

$$\oint_{\gamma} f = 0 \quad \forall \text{ great circles } \gamma$$

Funk 1914

$$\oint_{\gamma} f = 0 \quad \forall \gamma \iff f \text{ is odd} \quad (\text{cf. } \underline{\text{Radon 1917}})$$

Expect $\left\{ \begin{array}{l} S^2 \text{ is Blaschke } \underline{\text{deformable}} \quad (\checkmark \text{ Guillemin 1976}) \\ \mathbb{RP}_2 \text{ is Blaschke } \underline{\text{rigid}} \quad (\checkmark \dots \text{ LeBrun–Mason 2002}) \end{array} \right.$


Projective differential geometry

Defⁿ $\hat{\nabla}_a \sim \nabla_a \iff$ same geodesics (unparameterised)

EG (Thales 600 BC) the round sphere is projectively flat

Affine coordinate patch

$\mathbb{R}^n \hookrightarrow \mathbb{RP}_n$ is a
projective equivalence



Operational Defⁿ

$$\hat{\nabla}_a \phi_b = \nabla_a \phi_b - \Upsilon_a \phi_b - \Upsilon_b \phi_a$$

Projective deformations

Projective equivalence

$$\hat{\nabla}_a X^c = \nabla_a X^c + \Gamma_{ab}{}^c X^b \quad \text{where } \Gamma_{ab}{}^c = \Upsilon_a \delta_b^c + \Upsilon_b \delta_a^c$$

Projective deformation

$$\tilde{\nabla}_a X^c = \nabla_a X^c + \epsilon \Gamma_{ab}{}^c X^b \quad \text{where } \Gamma_{ab}{}^c = \Gamma_{(ab)}{}^c \text{ and } \Gamma_{ab}{}^a = 0$$

Projective deformation complex on S^n or \mathbb{RP}_n

$$\begin{array}{ccc} X^a & \mapsto & (\nabla_{(a} \nabla_{b)} X^c + g_{ab} X^c)_o & W_{abc}{}^d & \mapsto & \dots \\ & & \Gamma_{ab}{}^c & \mapsto & (\nabla_{[a} \Gamma_{b]c}{}^d)_o & \end{array}$$

(where $R_{abc}{}^d = g_{ac} \delta_b^d - g_{bc} \delta_a^d$ (round metric))

Riemannian deformations

Start with the round metric $R_{abcd} = g_{ac}g_{bd} - g_{bc}g_{ad}$

Recall Riemannian deformation

$$\tilde{g}_{ab} = g_{ab} + \epsilon h_{ab}$$

Riemannian deformation complex on S^n or $\mathbb{R}P_n$

$$X_a \mapsto \nabla_{(a} X_{b)} \qquad R_{abcd} \mapsto \nabla_{[a} R_{bc]de}$$

Killing

$$\begin{aligned} h_{ab} \mapsto & (\nabla_{(a} \nabla_{c)} + g_{ac}) h_{bd} \\ & - (\nabla_{(b} \nabla_{c)} + g_{bc}) h_{ad} \\ & - (\nabla_{(a} \nabla_{d)} + g_{ad}) h_{bc} \\ & + (\nabla_{(b} \nabla_{d)} + g_{bd}) h_{ac} \end{aligned}$$


Bianchi

Riemann tensor symmetries are SL-irreducible !


Projectively invariant complex !

Homogeneous correspondence


Homogeneous
under action
of $SL(n + 1, \mathbb{R})$



Hieroglyphics



$n = 3$



Homogeneous vector bundles



Differential Complexes on $\mathbb{R}P_3$

de Rham

$$0 \rightarrow \begin{array}{ccc} 0 & 0 & 0 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -2 & 1 & 0 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -3 & 0 & 1 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -4 & 0 & 0 \\ \times & \bullet & \bullet \end{array} \rightarrow 0$$

Riemannian deformation

$$0 \rightarrow \begin{array}{ccc} 0 & 1 & 0 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -2 & 2 & 0 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla^2} \begin{array}{ccc} -4 & 0 & 2 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -5 & 0 & 1 \\ \times & \bullet & \bullet \end{array} \rightarrow 0$$

Projective deformation

$$0 \rightarrow \begin{array}{ccc} 1 & 0 & 1 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla^2} \begin{array}{ccc} -3 & 2 & 1 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla} \begin{array}{ccc} -4 & 1 & 2 \\ \times & \bullet & \bullet \end{array} \xrightarrow{\nabla^2} \begin{array}{ccc} -6 & 1 & 0 \\ \times & \bullet & \bullet \end{array} \rightarrow 0$$

Generalised Funk transform on \mathbb{RP}_3

General complex

$$0 \rightarrow \begin{array}{c} a \quad b \quad c \\ \times \text{---} \bullet \text{---} \bullet \end{array} \xrightarrow{\nabla^{a+1}} \begin{array}{c} \text{\scriptsize } -a-2 \quad \text{\scriptsize } a+b+1 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \rightarrow \begin{array}{c} \text{\scriptsize } -a-b-3 \quad \text{\scriptsize } b+c+1 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \rightarrow \begin{array}{c} \text{\scriptsize } -a-b-c-4 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \rightarrow 0$$

$$\Gamma(\mathbb{RP}_3, \begin{array}{c} \text{\scriptsize } -a-2 \quad \text{\scriptsize } a+b+1 \\ \times \text{---} \bullet \text{---} \bullet \end{array}) \ni f \xrightarrow{\mathcal{F}} \oint_{\gamma} f \in \tilde{\Gamma}(\text{Gr}_2(\mathbb{R}^4), \begin{array}{c} a \quad b \quad c \\ \bullet \text{---} \times \text{---} \bullet \end{array})$$

Theorem (Bailey–E 1997) $\ker \mathcal{F} = \nabla^{a+1} (\Gamma(\mathbb{RP}_3, \begin{array}{c} a \quad b \quad c \\ \times \text{---} \bullet \text{---} \bullet \end{array}))$

Examples $\begin{array}{c} -2 \quad 2 \quad 0 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \iff$ Metric rigidity

$\begin{array}{c} -3 \quad 2 \quad 1 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \iff$ projective rigidity

$\begin{array}{c} -2 \quad m \quad 0 \\ \times \text{---} \bullet \text{---} \bullet \end{array} \iff$ injectivity of I_m on symmetric solenoidal fields

Generalised Funk transform on $\mathbb{C}\mathbb{P}_n$

Warning ☠️⚡☠️⚡☠️ $\mathbb{C}\mathbb{P}_n$ is not projectively flat!

$$W_{ab}{}^c{}_d = 2J_{[a}{}^c\omega_{b]d} - 2\omega_{ab}J_d{}^c - \frac{6}{2n-1}\delta_{[a}{}^c g_{b]d}$$

$$\Gamma(\mathbb{C}\mathbb{P}_n, \odot^{m-1}\Lambda^1) \xrightarrow{\nabla} \Gamma(\mathbb{C}\mathbb{P}_n, \odot^m\Lambda^1) \ni f \xrightarrow{I_m} \oint_{\gamma} f$$

Theorem (Tsukamoto 1981) $\ker I_2 = \text{range } \nabla$

Theorem (E-Goldschmidt 2013) $\ker I_m = \text{range } \nabla$

Technique $\mathbb{R}\mathbb{P}_n \hookrightarrow \mathbb{C}\mathbb{P}_n$ is totally geodesic and ...

?Theorem? (earlier today) $\mathbb{C}\mathbb{P}_n$ is projectively rigid



THE END

THANK YOU