


Fluctuations of Random Convex Hulls

Pierre Calka Joe Yukich


Banff BIRS, May 28- June 2, 2023

Introduction

- K : smooth convex body in \mathbb{R}^d .
- K_n : convex hull of n i.i.d. uniform points in K .
- For $K := \mathbb{B}^2, n = 500$, we have:



- How does the boundary of K_n fluctuate? Parabolic global constraints.



- Boundary: ∂K_n **a random interface**. As n increases, new points appear, creating new facets which may subsume existing facets.

When $n \rightarrow \infty$ we seek:

- limit distribution of the area of a facet chosen at random; limit distribution of distance between boundary of K and a facet chosen at random,
- limit distribution of *maximal* facet distance and *maximal* facet area,
- distributional convergence of process of heights of convex hull boundary,
- distributional convergence of process of heights for dynamic two-parameter process.

- $K \subset \mathbb{R}^d$ smooth C^3 convex body; $\kappa(\cdot) :=$ Gauss curvature along ∂K ; $\kappa > 0$
- Facets of K_n : Simplices a.s.
- \mathcal{F}_n : Facet chosen at random from the facets of K_n .
- $\text{dist}(\mathcal{F}_n)$: distance between the hyperplane containing \mathcal{F}_n and nearest supporting hyperplane.
- When K is the unit ball, $\text{dist}(\mathcal{F}_n) := 1 - \text{height}(\mathcal{F}_n)$.

Convergence in distribution of height/distance

- **Thm** As $n \rightarrow \infty$, we have $\mathbb{P}(n^{\frac{2}{d+1}} \text{dist}(\mathcal{F}_n) \leq t) \rightarrow 1 - F_{CH(K)}^{\text{height}}(t)$,

where

$$F_{CH(K)}^{\text{height}}(t) = c_d \int_{\partial K} \kappa(z)^{\frac{1}{d+1}} \int_0^\infty e^{-v} \left(v + \frac{1}{\sqrt{\kappa(z)}} \frac{\kappa_{d-1}}{d+1} (2t)^{(d+1)/2} \right)^{\frac{d(d-1)}{d+1}} \cdot \exp\left(-\frac{\kappa_{d-1}}{d+1} \frac{1}{\sqrt{\kappa(z)}} (2t)^{(d+1)/2}\right) dv dz.$$

- **Particular case** $K = \mathbb{B}^2$:


$$\mathbb{P}\left(\frac{n(1 - \text{height}(\mathcal{F}_n))}{n^{\frac{1}{3}}} \geq t\right) \sim Ct \exp\left(-\frac{4\sqrt{2}}{3} t^{\frac{3}{2}}\right) \quad \text{when } t \rightarrow \infty.$$

- $\mathbb{P}(n^{\frac{d-1}{d+1}} \text{Vol}_{d-1}(\mathcal{F}_n) \leq t) \rightarrow 1 - F_{CH(K)}^{\text{vol}}(t)$.


Process convergence of height function; unit ball \mathbb{B}^d

Notation $K = \mathbb{B}^d$

$\mathbf{H}(x, n) :=$
 height of K_n in the direction $x \in \mathbb{S}^{d-1}$;
 $\mathbf{H}(x, n) = \|x\| \mathbf{H}\left(\frac{x}{\|x\|}, n\right), \quad x \in \mathbb{R}^d$



$\exp_r \mathbb{S}^{d-1} :=$
 exponential map
 at the north pole of $r\mathbb{S}^{d-1}$





Theorem. Fix $L > 0$. As $n \rightarrow \infty$

$$\left\{ \frac{n - \mathbf{H}(n^{d/(d+1)} \exp_{n^{1/(d+1)} \mathbb{S}^{d-1}}(v), n)}{n^{(d-1)/(d+1)}} \right\}_{v \in B_{d-1}(0, L)} \xrightarrow{\mathcal{D}} \text{Burgers' festoon.}$$

$\cdot d = 2 : \quad \frac{1}{3} \quad \frac{2}{3} \text{ scaling}$

Convergence of the height function in dimension $1 + 1$



- Down paraboloid with apex at $(x_0, h_0) \in \mathbb{R} \times \mathbb{R}^+$:

$$\Pi^\downarrow(x_0, h_0) := \left\{ (x, h) \in \mathbb{R} \times \mathbb{R}, h - h_0 \leq -\frac{|x - x_0|^2}{2} \right\}.$$

- \mathcal{P} : Poisson pt process on $\mathbb{R} \times \mathbb{R}^+$. Burgers' festoon process Φ is

$$\Phi(x) := \sup_{(x_0, h_0) \in \mathbb{R} \times \mathbb{R}^+, \Pi^\downarrow(x_0, h_0) \cap \mathcal{P} = \emptyset} \left(h_0 - \frac{|x - x_0|^2}{2} \right).$$

- The parabolic faces in Φ are the re-scaled asymptotic images of the facets of $K_n, n \rightarrow \infty$.

Comparison between the convex hull interface and KPZ

- $d = 2$: $\frac{1}{3}$ $\frac{2}{3}$ scaling, but the limit process (Burgers' festoon) contains no Airy process.
- The marginal radial fluctuations converge to a limit distribution which has right-sided Tracy-Widom like tails.
- A time coordinate is missing (we return to this later).

- **Maximal radial fluctuation** = $MRF(K_n)$ = maximal facet distance.
- **Theorem** $MRF(K_n)$ asymptotically follows a Gumbel law, i.e., there are constants $a_i := a_i(K)$, $i \in \{0, 1, 2, 3\}$, such that if

$$t_n(\tau, K) := n^{-\frac{2}{d+1}} [a_0(a_1 \log n + a_2 \log(\log n) + a_3 + \tau)]^{\frac{2}{d+1}},$$

then as $n \rightarrow \infty$ we have

$$\mathbb{P}(MRF(K_n) \leq t_n(\tau, K)) \rightarrow \exp(-e^{-\tau}), \quad \tau \in (-\infty, \infty).$$

- $d = 2$: Bräker, Hsing, Bingham (1998).

Extreme Values: Facet Volumes

- $MFV(K_n) :=$ maximal volume of facets in ∂K_n
- **Theorem.** $MFV(K_n)$ asymptotically follows a Gumbel law, i.e., there are constants $b_i := b_i(K)$, $i \in \{0, 1, 2, 3\}$, such that if

$$t_n(\tau, K) := n^{-\frac{d-1}{d+1}} [b_0(b_1 \log n + b_2 \log(\log n) + b_3 + \tau)]^{\frac{d-1}{d+1}},$$

then as $n \rightarrow \infty$ we have

$$\mathbb{P}(MFV(K_n) \leq t_n(\tau, K)) \rightarrow \exp(-e^{-\tau}), \quad \tau \in (-\infty, \infty).$$

Growth of fluctuations; $d \geq 2$


- K_n : convex hull of an i.i.d. uniform sample in K of size n .
- **Corollary** (growth of fluctuations, with log precision in $d = 2$).

$$MRF(nK_n) \stackrel{\text{a.s.}}{=} \Theta(n^{1/3}(\log n)^{2/3}); \quad MFV(nK_n) \stackrel{\text{a.s.}}{=} \Theta(n^{2/3}(\log n)^{1/3}).$$

- $\frac{1}{3}, \frac{2}{3}$ scaling.
- Hammond: Convex hull boundary belongs to the **'baby KPZ class'**.
Global parabolic constraints, but no local Brownian fluctuations.
- $MRF(nK_n) \stackrel{\text{a.s.}}{=} \Theta(n^{\chi(d)}(\log n)^{\frac{2}{d+1}})$, $MFV(nK_n) \stackrel{\text{a.s.}}{=} \Theta(n^{\xi(d)}(\log n)^{\frac{1}{d+1}})$,
where $\chi(d) := (d-1)/(d+1)$ and $\xi(d) := d/(d+1)$ satisfy
 $\chi = 2\xi - 1$.
- Is there a two parameter space-time process?

Two parameter process: dynamic flower

- $X_i, 1 \leq i \leq n$, i.i.d. uniform in \mathbb{B}^2 . **Their flower** is $\bigcup_{i=1}^n B(\frac{X_i}{2}, \frac{|X_i|}{2})$:



Support function of **convex hull**
is boundary of flower, i.e.,

$$\max_{i \leq n} |X_i| \cos(|\theta - \theta_{X_i}|), \theta \in [0, 2\pi].$$

- Rescale radially by n , longitudinally by $\frac{1}{\sqrt{t}}$.
- 'Height of boundary of rescaled flower' at spatial coordinate θ at time $t > 0$:

$$h_n(\theta, t) = \max_{i \leq n} n |X_i| \cos\left(\frac{|\theta - \theta_{X_i}|}{\sqrt{t}\sqrt{n}}\right)$$

- For fixed n and large t the petals have nearly slope dependent growth.

Two parameter process: dynamic flower

- Re-scale space by $n^{2/3}$ and time by n :

$$h_n(n^{2/3}\theta, nt) = \max_{i \leq n} n|X_i| \cos\left(\frac{|\theta - \theta_{X_i}|}{\sqrt{t} \cdot n^{1/3}}\right).$$

- Define two parameter process with 1 : 2 : 3 scaling:

$$H_n(\theta, t) = \frac{h_n(n^{2/3}\theta, nt) - n}{n^{1/3}}, \quad \theta \in \mathbb{R}, t > 0.$$

- **Theorem.** Fix $t, L \geq 0$. As $n \rightarrow \infty$

$$\{H_n(\theta, t)\}_{|\theta| \leq L} \xrightarrow{\mathcal{D}} \{H(\theta, t)\}_{|\theta| \leq L}.$$

- Limit process given by variational formula (Burgers' festoon)

$$H(\theta, t) := \sup_{(v_0, h_0) \in \mathcal{P}} \left(h_0 - \frac{|\theta - v_0|^2}{2t} \right), \quad \theta \in \mathbb{R},$$

with \mathcal{P} a PPP in $\mathbb{R} \times \mathbb{R}_-$

- * Convergence in the space of cont. fcts on $|\theta| \leq L$ w. sup norm metric.

Two parameter process: dynamic flower

- **Theorem.** Fix $t, L \geq 0$. As $n \rightarrow \infty$

$$\{H_n(\theta, t)\}_{|\theta| \leq L} \xrightarrow{\mathcal{D}} \{H(\theta, t)\}_{|\theta| \leq L},$$

with

$$H(\theta, t) := \sup_{(v_0, h_0) \in \mathcal{P}} \left(h_0 - \frac{|\theta - v_0|^2}{2t} \right), \quad \theta \in \mathbb{R},$$

with \mathcal{P} a PPP in $\mathbb{R} \times \mathbb{R}_+$

- $\mathbb{P}(H_n(\theta_0, 1) \geq s) \rightarrow \exp(-\frac{4\sqrt{2}}{3}s^{3/2})$, $n \rightarrow \infty$.
- Shape profile is a pattern of coarsening paraboloids.
- $H_n(\theta, t)$ is an example of a process which satisfies 1:2:3 scaling, but does not belong to the KPZ fixed point universality class of Matetski, Quastel and Remenik (no Airy process).

Proof ideas: $d = 2$

- Fix n, t . Define the parabolic scaling transform $T^{(n)} : n\mathbb{B}^2 \rightarrow \mathbb{R} \times \mathbb{R}_-$

$$T^{(n)}(x) = \left(n^{-2/3} \exp_{n\mathbb{S}}^{-1}\left(\frac{x}{|x|}\right), n^{-1/3}(|x| - n) \right), \quad x \in n\mathbb{B}^2.$$

- $T^{(n)}$ maps $n\mathbb{S}^1$ to $[-\pi n^{1/3}, \pi n^{1/3}]$ and maps boundary of flower to a piecewise quasi-parabolic process $H^{(n)}$ in $[-\pi n^{1/3}, \pi n^{1/3}] \times [-n^{2/3}, 0]$.
- the shape of the quasi-parabolas constituting $H^{(n)}$ depends on n through via $T^{(n)}$; their apices are the image of an i.i.d. sample in $n\mathbb{B}^2$ under $T^{(n)}$, here denoted $\mathcal{P}^{(n)}$.
- the quasi-parabolas in the finite-area rectangle $[-L, L] \times [-\ell, 0]$, converge uniformly to parabolas in $[-L, L] \times [-\ell, 0]$, $n \rightarrow \infty$.
- couple $\mathcal{P}^{(n)}$ with a rate one Poisson point process on $\mathbb{R} \times \mathbb{R}_-$ such that with high probability they coincide on $[-L, L] \times [-\ell, 0]$.

Summary

- Limit distribution of height function of convex hull boundary has right-sided Tracy-Widom like tails in $d = 2$,
- Limit distributions of *maximal* facet distance and *maximal* facet area are of Gumbel type,
- Height process of convex hull boundary converges to Burgers' festoon,
- Height process of dynamic two-parameter flower converges to dynamic Burgers' festoon, with $1 : 2 : 3$ scaling and right-sided Tracy-Widom like tails in $d = 2$.

Thank you for your attention!